arXiv:1304.4723v1 [astro-ph.CO] 17 Apr 2013

Unavoidable strong magnetic fields in the early Universe

Jacques M. Wagstaff,[1] Robi Banerjee,[1] Dominik Schleicher,[2] and Günter Sigl[3]
[1] *Hamburger Sternwarte, Gojenbergsweg 112, 21029 Hamburg, Germany*[*]
[2] *Institut für Astrophysik, Georg-August-Universität Göttingen,*
Friedrich-Hund-Platz 1, 37077 Göttingen, Germany
[3] *II Institut für Theoretische Physik, Universität Hamburg,*
Luruper Chaussee 149, 22761 Hamburg, Germany

(Dated: April 18, 2013)

In this letter we show that the Universe is already strongly magnetised at very early epochs during cosmic evolution. Our calculations are based on the efficient amplification of weak magnetic seed fields, which are unavoidably present in the early Universe, by the turbulent small-scale dynamo (SSD). We identify at least one epoch during the radiation dominated regime where all the necessities for the SSD to work are fulfilled. Hence, at scales of $l_c \sim 0.3$ pc, the comoving field strength due to this mechanism will be $B_0 \sim 0.35\varepsilon^{1/2}$ nG at the present time, where ε is the saturation efficiency.

INTRODUCTION

Magnetic fields of strengths of order a few μG have been observed in high redshift as well as present day galaxies, and in galaxy clusters and superclusters (see e.g. refs. [1–4] respectively). There is also evidence for strong extragalactic magnetic fields coming from γ-ray observations. These studies place a lower bound on intergalactic magnetic fields at 3×10^{-16} G [5], although plasma effects may complicate the propagation of electromagnetic cascades [6, 7]. Theoretically, magnetic fields are very likely to have been generated at some level in the early Universe through a variety of mechanisms. On the largest scales, magnetic fields can be generated during inflation giving today $B_0 \sim 10^{-25} - 10^{-1}$ nG on a scale of 1 Mpc [8]. However such mechanisms require some modification to the Maxwell theory in order to break its conformal invariance. Without such modifications, fields of present strengths $B_0 \sim 10^{-20}$ and $\sim 10^{-11}$ nG could have been generated at the electroweak and QCD phase transitions respectively [9]. At later times, magnetic fields could have been generated through the generation of vorticity in the plasma of the early Universe [10–13] (originally proposed by Harrison [14]). The mechanism in refs. [10, 11] is very natural, since vorticity in the plasma is unavoidably generated in the late radiation era through the non-linear couplings of first order density perturbations. The seed field generated here is $B_0 \sim 10^{-20}$ nG. Therefore, in order to explain observations, some amplification of the seed fields must have occurred at some point in the history of the Universe. A popular mechanism for such amplification is known as the dynamo mechanism.

The dynamo mechanism has two classes (see ref. [15] for a review). The large-scale dynamo converts kinetic energy on large scales into magnetic energy. This mechanism can act if the conducting fluid flow is highly helical, inhomogeneous or anisotropic, the typical example being the differential rotation of galaxies. Whereas the SSD, which operates on much smaller scales, converts kinetic energy from turbulent motion into magnetic energy. Magnetic field lines which are frozen in the conducting plasma will be stretched by the random motion of fluid elements leading to field amplification. The SSD can take effect for nonhelical, homogeneous and isotropic turbulent flow. This mechanism has so far only been implemented to the formation of the first stars and galaxies in the matter dominated Universe [16] (see ref. [17] for an early discussion on this subject). In this paper we investigate the SSD amplification of a seed field in the radiation era. We demonstrate that the conditions are right for efficient dynamo amplification leading to large magnetic fields which could explain observations and have an impact on early structure formation.

TURBULENT FLUID VELOCITY IN THE RADIATION ERA

To explain the formation of the Large Scale Structure (LSS) in the Universe observed today, a primordial density perturbation of magnitude $\delta\rho/\rho \sim 10^{-5}$ is required at the time of matter-radiation equality. The primordial density perturbation is thought to have been generated at a much earlier time and therefore must be present in the very early Universe during the radiation dominated (RD) era. Cosmic inflation provides the most compelling explanation for the origin of the primordial curvature perturbation. A period of quasi-de Sitter inflation generically predicts a spatially flat Universe overlaid with a primordial spectrum of adiabatic, Gaussian and (nearly) scale-invariant superhorizon perturbations. Inflation generates scalar (density) and tensor (gravitational waves) modes. Vector modes, at linear level, are not sourced by inflation. The theory is in excellent agreement with observations of the LSS and CMB anisotropies [18, 19].

Well before horizon entry, the primordial curvature perturbation, which determines the gravitational potential Φ, remains constant and given by the initial condition Φ_0. In the Newtonian gauge, the equation of motion for the gravitational potential is given by [20]

$$\Phi'' + 3\mathcal{H}(1+w)\Phi' - w\nabla^2\Phi = 0, \tag{1}$$

where $p = w\rho$, $\mathcal{H} = a'/a$ and a prime denotes differentiation with respect to conformal time η. In the RD era $w = 1/3$, the solution to the above for the Fourier modes reads

$$\Phi(\boldsymbol{k},\eta) = \frac{3j_1(y)}{y}\Phi_0(\boldsymbol{k}), \tag{2}$$

where $y \equiv k\eta/\sqrt{3}$ and $j_1(y) = \sin y/y^2 - \cos y/y$ is the first spherical Bessel function. As a Fourier mode of the gravitational potential re-enters the horizon during RD it begins to oscillate with an amplitude decreasing as $1/y^2$.

The initial conditions, which are probed by observations, are given by the two-point correlation function

$$\langle\Phi_0(\boldsymbol{k}_1)\Phi_0^*(\boldsymbol{k}_2)\rangle = (2\pi)^3 P_\Phi(k)\delta^3(\boldsymbol{k}_1 - \boldsymbol{k}_2), \tag{3}$$

where

$$P_\Phi(k) = \frac{2\pi^2}{k^3}\frac{9}{25}\Delta^2_{\mathcal{R}}(k_0)\left(\frac{k}{k_0}\right)^{n_s-1}. \tag{4}$$

From the Planck results $\Delta^2_{\mathcal{R}}(k_0) = 2.215 \times 10^{-9}$ and $n_s \simeq 0.96$ for the pivot scale $k_0 = 0.05$ Mpc^{-1} [19]. Models of inflation are constructed in order to generate this amplitude and slight scale-dependence.

Perturbations in the fluid 3-velocity field are generated by the density perturbations. At first order in density perturbations, the fluid velocity perturbation is purely irrotational (curl-free) i.e. compressive turbulence, whose Fourier modes are [20]

$$v_i(\boldsymbol{k},\eta) = -\frac{ik_i}{2\mathcal{H}^2}\left[\Phi'(\boldsymbol{k},\eta) + \mathcal{H}\Phi(\boldsymbol{k},\eta)\right]. \tag{5}$$

With the solution in eq. (2) we find

$$v_i(\boldsymbol{k},\eta) = -\frac{i3\sqrt{3}\hat{k}_i}{2}\left[\sin y - 2j_1(y)\right]\Phi_0(k). \tag{6}$$

The perturbation in the velocity field oscillates with the density perturbation, however it has a term which does not decay with the expansion. The spectrum of velocity perturbations can be defined by the two-point correlation function

$$\langle v_i(\boldsymbol{k}_1,\eta)v_i^*(\boldsymbol{k}_2,\eta)\rangle = (2\pi)^3 P_v(k)\delta^3(\boldsymbol{k}_1 - \boldsymbol{k}_2), \tag{7}$$

where $\mathcal{P}_v(k) \equiv (k^3/2\pi^2)P_v(k)$. Considering eq. (6) we find

$$\langle v_i(\boldsymbol{k},\eta)v_i^*(\boldsymbol{k},\eta)\rangle = \frac{27}{4}\left[\sin y - 2j_1(y)\right]^2\langle|\Phi_0(k)|^2\rangle, \tag{8}$$

therefore

$$\mathcal{P}_v(k) = \frac{243}{100}\left[\sin y - 2j_1(y)\right]^2\Delta^2_{\mathcal{R}}(k_0)\left(\frac{k}{k_0}\right)^{n_s-1}. \tag{9}$$

For subhorizon scales $y, k/\mathcal{H} \gg 1$ the spectrum oscillates rapidly, so we can average the spectrum over many oscillations. For a scale-invariant primordial spectrum $n_s = 1$ we find

$$\overline{\mathcal{P}}_v(k) \simeq \frac{243}{200}\Delta^2_{\mathcal{R}}(k_0). \tag{10}$$

Thus, on subhorizon scales the spectrum of velocity perturbations becomes scale-invariant. The typical value of the velocity perturbation is therefore

$$v \sim \sqrt{\overline{\mathcal{P}}_v(k)} \simeq \sqrt{\Delta^2_{\mathcal{R}}(k_0)} \simeq 5 \times 10^{-5}, \tag{11}$$

where natural units are used such that $c = 1$. Hence, the turbulent velocity spectrum generated by first order density perturbations is isotropic, homogeneous, Gaussian and to a good approximation scale-invariant $v(l_p) \propto l_p^{\vartheta}$ with $\vartheta = 0$ on a physical scale l_p.



FIG. 1. The spectrum of velocity perturbations at temperature $T = T_f \simeq 19.6$ keV. The comoving length $l_c \simeq 0.27$ pc (black-dashed) is well within the oscillatory regime. (Color online)

THE SMALL-SCALE DYNAMO

In the SSD mechanism, kinetic energy from turbulent motion is converted to magnetic energy. There is a large number of numerical works in the literature which have demonstrated this effect unambiguously for a number of settings [21–24]. Generically, the fluctuating component of the magnetic field grows exponentially $B \propto \exp(\Gamma t)$, where the growth rate Γ depends on the kinetic Reynolds number R_e. Analytically, the Kazantsev theory (following the formalism by refs. [15, 25]) was developed in order to study the evolution of magnetic fields in a conducting plasma containing turbulent motion [26]. For random motions correlated on a scale l_p with velocity $v \propto l_p^{\vartheta}$, under certain conditions it can be shown that the growth rate is given by [27]

$$\Gamma = \frac{(163 - 304\vartheta)v}{60 l_p} R_e^{(1-\vartheta)/(1+\vartheta)}. \qquad (12)$$

This result is obtained in the large Prandtl number $P_m \equiv R_m/R_e$ limit, where the Prandtl number is a measure of the relative importance of the magnetic and kinetic diffusion. Here, the R_m is the magnetic Reynolds number. We note that the above analytical result was obtained by modelling the turbulent velocity spectrum so that it behaves as Kolmogorov or Burgers turbulence for scaling index $\vartheta = 1/3, 1/2$ respectively [27] (see conclusions for further discussion).

There is a critical magnetic Reynolds number for which $R_m > R_m^{\rm cr}$ and $\Gamma > 0$, and the SSD takes effect. The magnetic Reynolds number is given by $R_m = 4\pi l_p v\sigma$ where σ is the plasma conductivity. The local kinetic Reynolds number R_e characterises the relative importance of the fluid advective and dissipative terms in the Euler equation. On some physical length scale l_p, it is given by

$$R_e(l_p) = \frac{v(l_p) l_p}{\eta_s} = \frac{v(l_p) l_c}{(g_{\nu,\gamma}/5g_*) l_c^{\nu,\gamma}}, \qquad (13)$$

where l_c is a comoving length. The shear viscosity $\eta_s = (g_{\nu,\gamma}/5g_*) l_p^{\nu,\gamma}$ is determined by the particles of longest mean free path (*mfp*) $l_p^{\nu,\gamma}$, which are either neutrinos or photons in the radiation dominated era. Here, g_* and $g_{\nu,\gamma}$ are the total and component number of effective relativistic degrees of freedom. On a given scale l_p, a viscous regime corresponds to $R_e(l_p) \ll 1$. Turbulent flow corresponds to $R_e(l_p) \gg 1$ where the dissipative time-scale is much greater than the eddy-turnover time-scale $\tau_{\rm eddy} = l_p/v = a l_c/v$. We can now define the number of eddy-turnover times $N_{\rm eddy}$ given by integrating $\mathrm{d}N_{\rm eddy} = \mathrm{d}t/\tau_{\rm eddy}$. Since the epoch we are considering is in the RD era we have $a = T_0/T \propto t^{1/2}$ and

$$H = \pi \sqrt{\frac{g_*}{90}} \frac{T^2}{m_P} = \frac{1}{2t}, \qquad (14)$$

where we normalised $a_0 = 1$. The radiation temperature today is given by the CMB temperature $T_0 = 2.725$ K. With a constant v, given in eq. (11), we can integrate for $N_{\rm eddy}$ from some initial epoch defined by a temperature T_i, thus

$$N_{\rm eddy}(T, l_c) = \frac{v m_P}{\pi l_c T_0} \sqrt{\frac{90}{g_*}} \left(\frac{T_i - T}{T_i T} \right). \qquad (15)$$

The number of eddy-turnover times $N_{\rm eddy} \gtrsim 1$ so that the magnetic field amplification is not suppressed.

In the early Universe, before neutrino decoupling, neutrinos have the longest *mfp* and are thus most efficient at transporting momentum and heat. At high temperatures the shear viscosity due to neutrinos is low and the plasma is in a turbulent regime. However, the neutrino *mfp* increases as the Universe expands and cools, leading to a viscous regime. Eventually the neutrinos decouple at $T \simeq 2.6$ MeV and the fluid flow becomes turbulent once again since now the momentum is transported by photons which have small shear viscosity at this time. Once the temperature decreases below the electron mass $T < m_e = 0.511$ MeV, $e^\pm$ pairs begin to annihilate and the photon *mfp* increases rapidly. The $e^\pm$ annihilation completes at around $T \lesssim 20$ keV. After this, as the temperature drops further, photons begin to diffuse followed by photon drag and the fluid is in a viscous regime once again [28, 29]. The diffusion will also damp away the velocity fluctuations, see for example ref. [30]. Therefore, before diffusion sets in, there is always a turbulent regime if we consider large enough scales. However, we note that the eddy-turnover time increases for larger scales.

Let us consider the epoch of $e^\pm$ annihilation so that the initial temperature is $T_i = m_e$. At this time, at temperatures ranging $m_e \gtrsim T \gtrsim 20$ keV, the comoving photon *mfp* is given by [31]

$$l_c^\gamma = a^{-1} l_p^\gamma \simeq \frac{a^{-1}}{\sigma_T (n_{\rm pair}^2 + n_e^2)^{1/2}} \simeq \frac{a^{-1}}{\sigma_T n_{\rm pair}}, \qquad (16)$$

where $\sigma_T = 8\pi\alpha^2/3m_e^2$ is the Thomson cross section, $\alpha \approx 1/137$ is the fine structure constant and $n_{\rm pair}$ the number density of $e^\pm$ pairs given by [31]

$$n_{\rm pair} \approx \left(\frac{2 m_e T}{\pi} \right)^{3/2} \exp\left(-\frac{m_e}{T} \right) \left(1 + \frac{15}{8} \frac{T}{m_e} \right). \qquad (17)$$

At lower temperatures $l_c^\gamma \simeq a^{-1}/\sigma_T n_e$, where the number density of free electrons is

$$n_e = X_e n_b = X_e \frac{\rho_0}{m_p} \Omega_b \left(\frac{T}{T_0} \right)^3, \qquad (18)$$

and m_p is the proton mass. Since the plasma is fully ionised at this temperature the ionization fraction is $X_e = 1$.

The SSD amplification is efficient, i.e. not suppressed, for scales which have at least one eddy-turnover time whilst the kinetic Reynolds number is greater than unity. Let us define the final temperature T_f where turbulence is shut off by setting $R_e(T = T_f) = 1$ [32], i.e.

$$R_e(T = T_f) = \frac{v l_c \sigma_T T_0}{(g_\gamma/5g_*)} \frac{n_{\rm pair}(T_f)}{T_f} = 1. \qquad (19)$$

We can rearrange eq. (15) for $T = T_f$ and substitute it into eq. (19), from this we can find the largest possible scale l_c with $N_{\rm eddy}(T_f) \simeq 1$. With $g_* \simeq 3.36$, for the temperature range considered and the typical velocity fluctuations in eq. (11), we find the scale $l_c \simeq 0.27$ pc. This scale corresponds to $R_e(T_i) \sim 10^{12}$ and $T_f \simeq 19.6$ keV. This final temperature is close enough to the temperature at the end of $e^{\pm}$ completion to justify the approximation in eq. (16). Figure 1 shows the velocity spectrum at the final epoch T_f, the scale l_c above is clearly well inside the horizon where eq. (11) applies.

The magnetic Reynolds number is $R_m = 4\pi a l_c v\sigma$, where the conductivity at $T < m_e$ is given by [33]

$$\sigma \simeq \frac{m_e}{\alpha \ln \Lambda}\left(\frac{2T}{\pi m_e}\right)^{3/2}, \qquad (20)$$

and $\Lambda = (1/6\sqrt{\pi\alpha})\sqrt{m_e^3/n_e}(T/m_e)$. We can estimate the minimum conductivity in the epoch considered, which occurs at $T = T_f$, and thus the minimum R_m. With eq. (18) we find $R_m \sim 10^{15}$ for l_c and T_f given above, and thus the Prandtl number at this temperature is $P_m \equiv R_m/R_e \sim 10^{15}$. This large Prandtl number means that we can neglect dissipative effects due to finite conductivity. The large magnetic Reynolds number $R_m \gg R_m^{\rm cr} \sim \mathcal{O}(10^2)$ [15] justifies our choice of $R_e(T_f) = 1$ as the final state of efficient amplification.

For the length scale given above, the magnetic field growth rate Γ is much larger than the Hubble rate. From eq. (12), at the initial epoch considered, we find $\Gamma/H \sim 10^{11}$. Due to the exponential growth, the magnetic field will saturate very quickly within only one eddy-turnover time. Saturation is given by the equipartition between kinetic and magnetic energy. With no net magnetic helicity this occurs when [29]

$$\langle B^2\rangle \approx 4\pi\varepsilon(\rho+p)\langle v^2\rangle = \frac{16\pi^3\varepsilon}{90} g_* T^4 \langle v^2\rangle, \qquad (21)$$

which we evaluate at T_f and ε is the saturation efficiency (see discussion below). We find, with the kinetic energy given by the typical velocity fluctuations in eq. (11), $B \approx 3.7\varepsilon^{1/2} \times 10^6$ G, whose value today is $B_0 = B(T_0/T_f)^2 = 0.35\varepsilon^{1/2}$ nG. Although those strong fields are generated at an early epoch in the evolution of the Universe, they will not suffer further dynamical damping due to the flat velocity spectra and the survival of overdamped Alfven waves in the cosmic plasma [28, 29, 34]. The generation of a tiny seed field together with this efficient amplification via the SSD mechanism can easily explain the observed large extragalactic magnetic fields [5] and have a significant effect on a number of astrophysical processes.

DISCUSSION AND CONCLUSION

As noted in a number of numerical and analytical works, rapid amplification of seed magnetic fields occurs due to the turbulent motion of the conducting plasma. This small-scale dynamo (SSD) mechanism is believed to play a crucial role in the formation of large magnetic fields in a number of astronomical settings, from stars to galaxies and the intergalactic medium [16, 21, 22]. In this paper we have demonstrated that the conditions necessary for such turbulent amplification arise in the radiation era before the onset of structure formation. Turbulent plasma motion arise inevitably from perturbations of the gravitational potential; required for structure formation and believed to originate from quantum fluctuations of suitable fields during a period of cosmic inflation. Seed magnetic fields will almost certainly be generated at some level in the early Universe through a variety of mechanisms. Such mechanisms include inflation and phase transitions. A particularly natural mechanism, proposed by Harrison [14] and implemented in refs. [10–13], makes use of the vorticity generated by density perturbations and inevitably leads to the generation of seed fields. The generation of seed fields together with the SSD mechanism for their amplification could explain the observed large magnetic fields throughout the Universe.

The SSD mechanism is very efficient for rotational velocity fluctuations, where numerical simulations have shown that the saturation efficiency ε is close to unity [24]. The saturation level is lower for irrotational velocity fluctuations $\varepsilon \sim 10^{-3} - 10^{-4}$ [24], as we consider here. However, these numerical simulations were performed in a non-relativistic setting $c_s \ll 1$ and at relatively high Mach numbers. The required further numerical work to investigate the saturation level at low Mach numbers is left for future publications.

Rotational fluid velocity fluctuations $v_\perp$ are generated at second order in cosmological perturbations. Indeed, the generation of vector modes by density perturbations at second order, which contribute to the fluid 3-velocity, has been investigated by a number of works [35–38]. In ref. [36], the spectrum of vector modes was calculated for the radiation era. The power spectrum $\mathcal{P}_\mathcal{V}/\Delta^4_\mathcal{R}$ grows as y on superhorizon scales, peaks when inside the horizon and then decays as y^{-4} [36]. When the power spectrum peaks $\mathcal{P}_\mathcal{V} \sim \Delta^4_\mathcal{R}$. Hence, the maximum rotational velocity fluctuation is $v_\perp \sim \sqrt{\mathcal{P}_\mathcal{V}} \sim 10^{-9}$ and could contribute efficiently to the SSD action. However, the vector modes have a χ^2-distribution [36]. It is not clear how this non-Gaussianity affects the SSD action, further work in this direction is needed (although authors in ref. [39] suggest that breaking the Gaussian assumption does not change the qualitative results). Observational constraints however allow much greater amount of vector mode. In ref. [37], neglecting gravitational waves, the authors showed that $\mathcal{P}_\mathcal{V}/\mathcal{P}_s \lesssim 0.012$ from WMAP 7-year data. If such large vector mode exists, the SSD amplification could be very efficient in the early Universe. Indeed, this opens up the intriguing possibility of efficient magnetic field amplification due to vector modes generated in non-standard models of inflation.

Finally, we stress that the main assumptions of the Kazantsev model are simplifying assumptions needed to obtain an analytical grip on the problem. These assumptions regard the statistical properties of the velocity field.

The velocity field is taken to be Gaussian, homogeneous and isotropic in space and instantaneously correlated in time. So far we have not yet calculated the detailed statistical properties of the velocity fluctuations. We have only seen that the irrotational component has a Gaussian distribution whereas the rotational component is non-Gaussian, and neither are instantaneously correlated in time (see for example ref. [40] for related numerical work). For this reason it would be of high interest to develop an analytical model for more general velocity fields. We leave this work for future publications. Nevertheless, we expect that below the integral scale of the irrotational velocity fluctuation spectrum Kolmogorov type turbulence is established. But we stress that the Kazantsev mechanism will work independently of the type of turbulence [41]. Hence the early strong magnetisation of the Universe seems unavoidable.

This work was supported by the Deutsche Forschungsgemeinschaft through the collaborative research centre SFB 676, by the *Helmholtz Alliance for Astroparticle Phyics* (HAP) funded by the Initiative and Networking Fund of the Helmholtz Association. DRGS thanks for funding by the German Science Foundation (DFG) via the SFB 963/1 *Astrophysical Flow Instabilities and Turbulence* (project A12).

---

* jwagstaff@hs.uni-hamburg.de

[1] R. Beck, Space Science Reviews **166**, 215 (2012).

[2] M. L. Bernet, F. Miniati, S. J. Lilly, P. P. Kronberg, and M. Dessauges-Zavadsky, Nature **454**, 302 (2008), arXiv:0807.3347.

[3] L. Feretti, G. Giovannini, F. Govoni, and M. Murgia, A&AR **20**, 54 (2012), arXiv:1205.1919 [astro-ph.CO].

[4] Y. Xu, P. P. Kronberg, S. Habib, and Q. W. Dufton, AstrophysJ **637**, 19 (2006), arXiv:astro-ph/0509826.

[5] A. Neronov and I. Vovk, Science **328**, 73 (2010), arXiv:1006.3504 [astro-ph.HE].

[6] A. E. Broderick, P. Chang, and C. Pfrommer, Astrophys. J. **752**, 22 (2012), arXiv:1106.5494 [astro-ph.CO].

[7] R. Schlickeiser, D. Ibscher, and M. Supsar, Astrophys. J. **758**, 102 (2012).

[8] M. S. Turner and L. M. Widrow, Phys.Rev. **D37**, 2743 (1988).

[9] G. Sigl, A. V. Olinto, and K. Jedamzik, Phys.Rev. **D55**, 4582 (1997), arXiv:astro-ph/9610201 [astro-ph].

[10] S. Matarrese, S. Mollerach, A. Notari, and A. Riotto, Phys.Rev. **D71**, 043502 (2005), arXiv:astro-ph/0410687 [astro-ph].

[11] K. Takahashi, K. Ichiki, H. Ohno, and H. Hanayama, Phys.Rev.Lett. **95**, 121301 (2005), arXiv:astro-ph/0502283 [astro-ph].

[12] E. Fenu, C. Pitrou, and R. Maartens, Mon.Not.Roy.Astron.Soc. **414**, 2354 (2011), arXiv:1012.2958 [astro-ph.CO].

[13] T. Kobayashi, R. Maartens, T. Shiromizu, and K. Takahashi, Phys.Rev. **D75**, 103501 (2007), arXiv:astro-ph/0701596 [astro-ph].

[14] E. Harrison, Mon.Not.Roy.Astron.Soc. **147**, 279 (1970).

[15] A. Brandenburg and K. Subramanian, Phys.Rept. **417**, 1 (2005), arXiv:astro-ph/0405052 [astro-ph].

[16] D. R. G. Schleicher, R. Banerjee, S. Sur, T. G. Arshakian, R. S. Klessen, R. Beck, and M. Spaans, A&A **522**, A115 (2010), arXiv:1003.1135 [astro-ph.CO].

[17] R. Beck, A. Brandenburg, D. Moss, A. Shukurov, and D. Sokoloff, ARA&A **34**, 155 (1996).

[18] G. Hinshaw, D. Larson, E. Komatsu, D. Spergel, C. Bennett, *et al.*, (2012), arXiv:1212.5226 [astro-ph.CO].

[19] P. Ade *et al.* (Planck Collaboration), (2013), arXiv:1303.5076 [astro-ph.CO].

[20] H. Kodama and M. Sasaki, Prog.Theor.Phys.Suppl. **78**, 1 (1984).

[21] M. Latif, D. Schleicher, W. Schmidt, and J. Niemeyer, (2012), arXiv:1212.1619 [astro-ph.CO].

[22] S. Sur, D. R. G. Schleicher, R. Banerjee, C. Federrath, and R. S. Klessen, Astrophys. J., Letters **721**, L134 (2010), arXiv:1008.3481 [astro-ph.CO].

[23] L. Haugen, A. Brandenburg, and W. Dobler, Astrophys.Space Sci. **292**, 53 (2004), arXiv:astro-ph/0306453 [astro-ph].

[24] C. Federrath, G. Chabrier, J. Schober, R. Banerjee, R. S. Klessen, *et al.*, Phys.Rev.Lett. **107**, 114504 (2011), arXiv:1109.1760 [physics.flu-dyn].

[25] K. Subramanian, (1997), arXiv:astro-ph/9708216 [astro-ph].

[26] A. Kazantsev, JHEP **53**, 1806 (1967).

[27] J. Schober, D. Schleicher, C. Federrath, R. Klessen, and R. Banerjee, Phys.Rev. E **85**, 026303 (2012), arXiv:1109.4571 [astro-ph.CO].

[28] K. Jedamzik, V. Katalinic, and A. V. Olinto, Phys.Rev. **D57**, 3264 (1998), arXiv:astro-ph/9606080 [astro-ph].

[29] R. Banerjee and K. Jedamzik, Phys.Rev. **D70**, 123003 (2004), arXiv:astro-ph/0410032 [astro-ph].

[30] S. Dodelson, *Modern cosmology* (Academic Press, 2003).

[31] K. Jedamzik and G. M. Fuller, Astrophys.J. **423**, 33 (1994), arXiv:astro-ph/9312063 [astro-ph].

[32] Any higher threshold for the Reynolds number will just slightly lower the coherence length.

[33] J. D. Jackson, *Classical Electrodynamics* (John Wiley and Sons, New York, 1975).

[34] K. Jedamzik and G. Sigl, Phys.Rev. **D83**, 103005 (2011), arXiv:1012.4794 [astro-ph.CO].

[35] T. H.-C. Lu, K. Ananda, C. Clarkson, and R. Maartens, JCAP **0902**, 023 (2009), arXiv:0812.1349 [astro-ph].

[36] T. H.-C. Lu, K. Ananda, and C. Clarkson, Phys.Rev. **D77**, 043523 (2008), arXiv:0709.1619 [astro-ph].

[37] K. Ichiki, K. Takahashi, and N. Sugiyama, Phys.Rev. **D85**, 043009 (2012), arXiv:1112.4705 [astro-ph.CO].

[38] A. Lewis, Phys.Rev. **D70**, 043518 (2004), arXiv:astro-ph/0403583 [astro-ph].

[39] S. Boldyrev and F. Cattaneo, Phys.Rev.Lett. **92**, 144501 (2004), arXiv:astro-ph/0310780 [astro-ph].

[40] J. Mason, L. Malyshkin, S. Boldyrev, and F. Cattaneo, Astrophys.J. **730**, 86 (2011), arXiv:1101.5181 [physics.plasm-ph].

[41] A. Gruzinov, S. C. Cowley, and R. Sudan, Phys.Rev.Lett. **77**, 4342 (1996), arXiv:astro-ph/9611194 [astro-ph].